

Recombinant Human Arginine decarboxylase

| | |
|---------------------------|---|
| Catalogue No.: | P6365 |
| Species: | Human |
| Uniprot ID: | Q96A70 |
| Expression Region: | 167-411 |
| Host: | E.Coli |
| Tags: | N-terminal His Tag or N-terminal His-IF2DI Tag, determined during production process |
| Molecular Weight: | TBD |
| Purity: | Greater than 95% as determined by SDS-PAGE |
| Formulation: | Lyophilized from a 0.2 μ m filtered solution in 10 mM Hepes, 500 mM NaCl with 5% trehalose, pH7.4 |
| Reconstitution: | Centrifuge the vial prior to opening, reconstitute in sterile distilled water or buffer of interest to a concentration of 0.1-1mg/ml by gently pipetting 2-3 times, don't vortex. |
| Storage: | The lyophilized protein is stable at -20°C for up to 1 year, the protein solution can be stored at 2-8°C for up to 1 week. For extended storage, it is recommended to further dilute in a buffer containing 0.1% BSA (make sure BSA will NOT interfere with your experiment) and store in working aliquots at -20°C to -80°C for 6 months. Avoid repeated freeze/thaw cycle. |

Synonyms: ADC, ADC_HUMAN, Antizyme inhibitor 2, ArgDC, Arginine decarboxylase, AZI2, KIAA1945, L-Arginine carboxy lyase, L-Arginine decarboxylase, ODC antizyme inhibitor 2, ODC paralog, ODC-p, ODC-paralogue, ODC1L, ODCp, Ornithine decarboxylase like, Ornithine decarboxylase paralog, Ornithine decarboxylase-like protein

SDS-PAGE: Test in progress

Safety Note: This product is intended for research and manufacturing uses only. It is not a diagnostic device. Product degradation will result from multiple freeze/thaw cycles. It is suggested that the antigen be stored in use size aliquots and thawed just prior to use. This material has been inactivated, however as with all biological materials, it should be handled as potentially infectious. The user assumes all responsibility for care, custody and control of the material, including its disposal, in accordance with all regulations.